



## G - กึ่งกรุปอันดับที่บรรจุฐานสองด้าน

### On Ordered $G$ - Semigroups Containing Two-sided Bases

วิชญาพร จันทะนัน, กชกร นวดไธสง และ สุชาติ เจริญรัมย์\*

Wichayaporn Jantanana, Kodchakorn Nuadthaisong และ Suchat Jaroenram\*

ภาควิชาคณิตศาสตร์ คณะวิทยาศาสตร์ มหาวิทยาลัยราชภัฏบุรีรัมย์

Department of Mathematics, Faculty of Science, Buriram Rajabhat University

Received : 2 September 2020

Revised : 15 December 2020

Accepted : 26 February 2021

#### บทคัดย่อ

ในบทความวิจัยนี้ศึกษาแนวคิดของ แกมมา กึ่งกรุปอันดับที่บรรจุฐานสองด้าน โดยจะศึกษาคัดลอกกับแนวคิดของ แกมมา กึ่งกรุปที่บรรจุฐานสองด้าน ซึ่ง ธวัช ช่างฝัด และ พิสิทธิ์ คำมูล ได้ทำการศึกษาในปี ค.ศ. 2018 นอกจากนี้ เรายังพิสูจน์ว่าทุก ๆ แกมมา กึ่งกรุปอันดับที่บรรจุฐานสองด้านจะมีสมาชิกเท่ากัน

คำสำคัญ : แกมมา กึ่งกรุปอันดับ ; ฐานสองด้าน ; แกมมาไอดีล

#### Abstract

The aim of this paper is to study the concept of ordered  $G$  - semigroups containing two-sided bases that are studied analogously to the concept of  $G$  - semigroups containing two-sided bases considered by T. Changpas and P. Kummoon in 2018. Moreover, we prove any ordered  $G$  - semigroups containing two-sided bases have the same cardinality.

Keywords : ordered  $G$  - semigroup ; two-sided bases ;  $G$  - Ideal

\*Corresponding author. E-mail : 600112210021@bru.ac.th



**Introduction**

The notion of two sided bases of semigroup has been introduced and studied by I. Fabrici. (Fabrici, 1975). Indeed, a non-empty subset  $A$  of semigroup  $S$  is said to be a two-sided bases of  $S$  if  $A$  satisfies the following two conditions :

$$(1) S = A \dot{\cup} SA \dot{\cup} AS \dot{\cup} SAS.$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = B \dot{\cup} SB \dot{\cup} BS \dot{\cup} SBS, \text{ then } B = A.$$

The concept of a  $G$ -semigroup has been introduced by M. K. Sen. (Sen, 1981). The concept of  $G$ -semigroup containing two sided bases was first given by T. Changpas and P. Kummoon. (Thawat & Pisit, 2018). Which form of  $G$ -semigroup containing two sided bases is a non-empty subset  $A$  of a  $G$ -semigroup  $S$  is called a two-sided bases of  $S$  it is satisfies the following two conditions :

$$(1) S = A \dot{\cup} SGA \dot{\cup} AGS \dot{\cup} SGA GS.$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = B \dot{\cup} SGB \dot{\cup} BGS \dot{\cup} SGB GS, \text{ then } B = A.$$

The main purpose of this paper is to introduce the concept and extend the result to on ordered  $G$ -semigroup containing two-sided bases. Will get the form of ordered  $G$ -semigroup containing two-sided bases is a non-empty subset  $A$  of an ordered  $G$ -semigroup  $S$  is called a two-sided bases of  $S$  it is satisfies the following two conditions:

$$(1) S = (A \dot{\cup} SGA \dot{\cup} AGS \dot{\cup} SGA GS).$$

$$(2) \text{ If } B \text{ is a subset of } A \text{ such that } S = (B \dot{\cup} SGB \dot{\cup} BGS \dot{\cup} SGB GS), \text{ then } B = A.$$

We now recall some definition and results used throughtout the paper.

**Definition 1.1.** (Thawat & Pisit, 2018). Let  $S$  and  $G$  be any two non-empty sets. Then  $S$  is called a  $G$ -semigroup if there exists a mapping from  $S \times G \times S \rightarrow S$ , written as  $(a, g, b) \mapsto agb$ , satisfying the following identity  $(aab)bc = aa(bbc)$  for all  $a, b, c \in S$  and  $a, b \in G$ .

**Definition 1.2.** (Abdul et al., 2017). Let  $(S, G, \leq)$  be an ordered  $G$ -semigroup. For  $A$  and  $B$  be two non-empty subsets of  $S$ , the set product  $AGB$  is defined to be the set of all elements  $agb$  in  $S$  where  $a \in A, b \in B$  and  $g \in G$ . That is

$$AGB := \{agb \mid a \in A, b \in B, g \in G\}.$$

Also we write  $BGa$  instead of  $BG\{a\}$ , and similarly for  $aGB$ , for  $a \in S$ .

**Definition 1.3.** (Niovi, 2017). An ordered  $G$ -semigroup is a  $G$ -semigroup  $S$  together with an order relation  $\leq$  such that  $a \leq b$  implies  $agc \leq bgc$  and  $cga \leq cgb$  for all  $a, b, c \in S$  and  $g \in G$ .

(lampan, 2009). For an element  $a$  of ordered  $G$ -semigroup  $S$ , define  $(a) := \{t \in S \mid t \leq a\}$  and for a subset  $H$  of  $S$ , define  $(H) = \bigcup_{h \in H} (h)$  that is  $(H) = \{t \in S \mid t \leq h \text{ for some } h \in H\}$ . Then following holds true:



1.  $H \cap (H) = (H) \cap H$ ;
2. For any subsets  $A$  and  $B$  of  $S$  with  $A \subseteq B$ , we have  $(A] \subseteq (B]$ ;
3. For any subsets  $A$  and  $B$  of  $S$ , we have  $(A \cap B] = (A] \cap (B]$ ;
4. For any subsets  $A$  and  $B$  of  $S$ , we have  $(A \cup B] \subseteq (A] \cup (B]$ .

**Definition 1.4.** (Niovi, 2017). A non-empty subset  $A$  of an ordered  $G$ -semigroup  $(S, G, \leq)$  is called a  $G$ -subsemigroup (or simply a subsemigroup) of  $S$  if  $AG \cap A = A$ .

**Definition 1.5.** (Kwon and Lee, 1998). A non-empty subset  $A$  of an ordered  $G$ -semigroup  $(S, G, \leq)$  is called a left (resp. right)  $G$ -ideal of  $S$  if it satisfies :

- (1)  $SG \cap A = A$  (resp.  $AG \cap A = A$ )
- (2) if  $a \in A$  and  $b \leq a$  for  $b \in S$  implies  $b \in A$ .

Both a left  $G$ -ideal and a right  $G$ -ideal of an ordered  $G$ -semigroup  $S$  is called a  $G$ -ideal of  $S$ .

**Definition 1.6.** (Kostaq & Edmond, 2006). An  $G$ -ideal  $A$  of an ordered  $G$ -semigroup  $(S, G, \leq)$  is called proper if  $A \neq S$ . A proper ideal  $A$  of  $S$  is called maximal if for each  $G$ -ideal  $T$  of  $S$  such that  $A \subseteq T$ , we have  $T = A$  or  $T = S$  i.e., there is no  $G$ -ideal  $T$  of  $S$  such that  $A \subsetneq T \subsetneq S$ .

**Proposition 1.7.** (Kostaq & Edmond, 2006). Let  $(S, G, \leq)$  be an ordered  $G$ -semigroup and  $\{A_i \mid i \in I\}$  a non-empty family of ideals of  $S$ . If  $\bigcap_{i \in I} A_i \neq \emptyset$ , then the set  $\bigcap_{i \in I} A_i$  is a  $G$ -ideal of  $S$  and  $\bigcup_{i \in I} A_i$  is also a  $G$ -ideal of  $S$ .

It is known (Niovi, 2017) that if denoted by  $I(A)$ , is the smallest  $G$ -ideal of  $S$  containing  $A$ , and  $I(A)$  is of the form  $I(A) = (A \cup SG \cup AG \cup SGA \cup SGA \cup GS]$ . In particular, for an element  $a \in S$ , we write  $I(\{a\})$ ,  $I(a)$  which is called the principal  $G$ -ideal of  $S$  generated by  $a$ . Thus  $I(a) = (a \cup SGa \cup aGS \cup SGA \cup SGA \cup GS]$ . Note that for any  $b \in S$ , we have  $(SGB \cup bGS \cup SGBGS]$  is a  $G$ -ideal of  $S$ . Finally, if  $A$  and  $B$  are two  $G$ -ideal of  $S$ , then the union  $A \cup B$  is a  $G$ -ideal of  $S$ .

## Methods

We begin this section with the definition of two-sided bases of ordered  $G$ -semigroup as follows.

**Definition 2.1.** (Abul *et al.*, 2017). Let  $(S, G, \leq)$  be an ordered  $G$ -semigroup. A non-empty subset  $A$  of  $S$  is called a two-sided base of  $S$  if it satisfies the following two conditions.

- (1)  $S = (A \cup SGA \cup AGS \cup SGA \cup GS]$ .
- (2) If  $B$  is a subset of  $A$  such that  $S = (B \cup SGB \cup BGS \cup SGBGS]$ , then  $B = A$ .

We now provide some examples.



**Example 2.2.** (Chinnadurai & Arulmozhi, 2018). Let  $S = \{a, b, c, d\}$  and  $G = \{a, b\}$  where  $a, b$  is defined on  $S$  with the following Cayley tables:

$a$	$a$	$b$	$c$	$d$
$a$	$a$	$a$	$a$	$a$
$b$	$a$	$b$	$c$	$d$
$c$	$a$	$c$	$c$	$c$
$d$	$a$	$c$	$c$	$c$

$b$	$a$	$b$	$c$	$d$
$a$	$a$	$a$	$a$	$a$
$b$	$a$	$b$	$c$	$d$
$c$	$a$	$c$	$c$	$c$
$d$	$a$	$b$	$c$	$d$

$$\mathcal{E} := \{(a, a), (a, b), (a, c), (a, d), (b, b), (b, c), (b, d), (c, c), (d, c), (d, d)\}$$

In (Chinnadurai & Arulmozhi, 2018).  $(S, G, \mathcal{E})$  is an ordered  $G$ -semigroup. It is easy to see that the two-sided bases of  $S$  are  $\{b\}$  and  $\{d\}$ . But  $\{b, d\}$  is not a two-sided base

**Example 2.3.** (Subrahmanyeswara *et al.*, 2012). Let  $S = \{a, b, c, d\}$  and  $G = \{a\}$  where  $a$  is defined on  $S$  with the following Cayley tables:

$a$	$a$	$b$	$c$	$d$
$a$	$b$	$b$	$d$	$d$
$b$	$b$	$b$	$d$	$d$
$c$	$d$	$d$	$c$	$d$
$d$	$d$	$d$	$d$	$d$

$$\mathcal{E} := \{(a, a), (b, b), (c, c), (d, d), (a, b), (d, b), (d, c)\}$$

In (Subrahmanyeswara *et al.*, 2012).  $(S, G, \mathcal{E})$  is an ordered  $G$ -semigroup. It is easy to see that the two-sided bases of  $S$  is  $\{a, c\}$ . But  $\{b\}$  and  $\{d\}$  is not a two-sided base

In Example 2.2. and 2.3., it is observed that two-sided bases of  $S$  have same cardinality. This leads to prove in Theorem 3.4.

Hereafter, for any ordered  $G$ -semigroup  $(S, G, \mathcal{E})$ , we shall use the quasi-ordering which is defined as follows.

**Definition 2.4.** Let  $(S, G, \mathcal{E})$  be an ordered  $G$ -semigroup. We define a quasi-ordering on  $S$  by for any  $a, b \in S$ ,

$$a \leq b \iff I(a) \subseteq I(b).$$

We write  $a \leq b$  if  $a \leq b$  but  $a \neq b$ . It is clear that, for any  $a, b \in S$ ,  $a \leq b$  implies  $a \leq b$ .

**Lemma 2.5.** Let  $A$  be a two-sided base of an ordered  $G$ -semigroup  $(S, G, \mathcal{E})$ , and  $a, b \in A$ . If  $a \leq (S G b \mathcal{E} b G S \mathcal{E} S G b G S)$ ,  $\mathcal{E} S G b G S$ , then  $a = b$ .



**Proof.** Assume that  $a \in (SGB \dot{=} bGS \dot{=} SGBGS]$ , and suppose that  $a \neq b$ . Let  $B = A \setminus \{a\}$ . Since  $a \neq b$ ,  $b \in B$ . To show that  $I(A) \dot{=} I(B)$ , it suffices to show that  $A \dot{=} I(B)$ . Let  $x \in A$ . There are two cases to consider. If  $x \neq a$ , then  $x \in B$ , and so  $x \in I(B)$ . If  $x = a$ , then by assumption we have  $x = a \in (SGB \dot{=} bGS \dot{=} SGBGS] \dot{=} I(b) \dot{=} I(B)$ . So we have  $I(A) \dot{=} I(B)$ . Thus  $S = I(A) \dot{=} I(B) \dot{=} S$ . This is contradiction. Hence  $a = b$ .

### Results

In this part the algebraic structure of an ordered  $G$ -semigroup containing two-sided bases will be presented.

**Theorem 3.1.** A non-empty subset  $A$  of an ordered  $G$ -semigroup  $(S, G, \leq)$  is a two-sided base of  $S$  if and only if  $A$  satisfies the following two conditions:

- (1) For any  $x \in S$  there exists  $a \in A$  such that  $x \leq_l a$ ;
- (2) For any  $a, b \in A$ , if  $a \neq b$ , then neither  $a \leq_l b$  nor  $b \leq_l a$ .

**Proof.** Assume first that  $A$  is a two-sided base of  $S$ . Then  $I(A) = S$ . Let  $x \in S$ . Then  $x \in I(A) = \{I(a) \mid \text{for all } a \in A\}$ , and so  $x \in I(a)$  for some  $a \in A$ . This implies  $I(x) \dot{=} I(a)$ . Hence  $x \leq_l a$ . Thus (1) holds. Let  $a, b$  be element of  $A$  such that  $a \neq b$ . Suppose  $a \leq_l b$ . We set  $B = A \setminus \{a\}$ . Then  $b \in B$ . Let  $x$  be element of  $S$ . By (1), there exists  $c$  in  $A$  such that  $x \leq_l c$ . There are two cases to consider. If  $c \neq a$ , then  $c \in B$ , thus  $I(x) \dot{=} I(c) \dot{=} I(B)$ . Hence  $S = I(B)$ . This is a contradiction. If  $c = a$ , then  $x \leq_l a$  hence  $x \in I(B)$  since  $b \in B$ . We have  $S = I(B)$ . This is a contradiction. The case  $b \leq_l a$  is proved similary. Thus (2) holds true.

Conversaly, assume that the condition (1) and (2) hold. We will show that  $A$  is a two-sided base of  $S$ . To show that  $S = I(A)$ . Let  $x \in S$ . By (1), Then there exists  $a \in A$  such that  $I(x) \dot{=} I(a)$ . Then  $x \in I(x) \dot{=} I(a) \dot{=} I(A)$ . Thus  $S \dot{=} I(A)$ , and  $S = I(A)$ . Next it remains to show that  $A$  is a minimal subset of  $S$  with the property:  $S = I(A)$ . Suppose that  $S = I(B)$  for some  $B \dot{=} A$ . Since  $B \dot{=} A$ , there exists  $a \in A \setminus B$ . So  $a \notin B$ . Since  $a \in A \dot{=} S = I(B)$  and  $a \notin B$ , it follows that  $a \in (SGB \dot{=} BGS \dot{=} SGBGS]$ . Since  $a \in (SGB \dot{=} BGS \dot{=} SGBGS]$ , we have  $a \leq y$  for some  $y \in SGB \dot{=} BGS \dot{=} SGBGS$ . There are three cases to consider:

**Case 1:**  $y \in BGS$ . Then  $y = b_1gs$  for some  $b_1 \in B, g \in G$  and  $s \in S$ . Since  $a \leq y$  and  $y \in b_1 \dot{=} SGB_1 \dot{=} b_1GS \dot{=} SGB_1GS$ . So  $a \in (b_1 \dot{=} SGB_1 \dot{=} b_1GS \dot{=} SGB_1GS]$ , it follows that  $I(a) \dot{=} I(b_1)$ . Hence,  $a \leq_l b_1$ . This is a contradiction.

**Case 2:**  $y \in SGB$ . Then  $y = sgb_2$  for some  $b_2 \in B, g \in G$  and  $s \in S$ . Since  $a \leq y$  and  $y \in b_2 \dot{=} SGB_2 \dot{=} b_2GS \dot{=} SGB_2GS$ . So  $a \in (b_2 \dot{=} SGB_2 \dot{=} b_2GS \dot{=} SGB_2GS]$ , it follows that  $I(a) \dot{=} I(b_2)$ . Hence,  $a \leq_l b_2$ . This is a contradiction.



**Case 3:**  $y \in SGBGS$ . Then  $y = s_1g_1b_3g_2s_2$  for some  $b_3 \in B, g_1, g_2 \in G$  and  $s \in S$ . Since  $a \notin y$  and  $y \in b_3 \in SGB_3 \in b_3GS \in SGB_3GS$ . So  $a \in (b_3 \in SGB_3 \in b_3GS \in SGB_3GS)$ , it follows that  $I(a) \in I(b_3)$ . Hence  $a \leq b_3$ . This is a contradiction.

Therefore  $A$  is a two-sided base of  $s$  as required, and the proof is completed.

**Theorem 3.2.** Let  $A$  be a two-sided base of an ordered  $G$ -semigroup  $(S, G, \leq)$  such that  $I(a) = I(b)$  for some  $a$  in  $A$  and  $b$  in  $S$ . If  $a \neq b$ , then  $S$  contains at least two-sided base.

**Proof.** Assume that  $a \neq b$ . Suppose that  $b \in A$ . Since  $a \leq b$  and  $a \neq b$ , it follows that  $a \in (SGB \in bGS \in SGBGS)$ . By Lemma 2.5., we obtain  $a = b$ . This is a contradiction. Thus  $b \in S \setminus A$ . Let  $B := (A \setminus \{a\}) \cup \{b\}$ . Since  $b \in B$ , we have  $b \in A$ , and  $B \not\subseteq A$ . Hence  $A \neq B$ . We will show that  $B$  is a two-sided base of  $S$ . To show that  $B$  satisfies (1) in Theorem 3.1., let  $x \in S$ . Since  $A$  is a two-sided base of  $S$ , there exists  $c \in A$  such that  $x \leq c$ . If  $c \neq a$ , then  $c \in B$ . If  $c = a$ , then  $x \leq a$ . Since  $a \leq b, x \leq a \leq b$ , we have  $x \leq b$ . To show that  $B$  satisfies (2) in Theorem 3.1., let  $c_1, c_2 \in B$  be such that  $c_1 \neq c_2$ . We will show that neither  $c_1 \leq c_2$  nor  $c_2 \leq c_1$ . Since  $c_1 \in B$  and  $c_2 \in B$ , we have  $c_1 \in A \setminus \{a\}$  or  $c_1 = b$  and  $c_2 \in A \setminus \{a\}$  or  $c_2 = b$ . There are four cases to consider:

**Case 1:**  $c_1 \in A \setminus \{a\}$  and  $c_2 \in A \setminus \{a\}$ . This implies neither  $c_1 \leq c_2$  nor  $c_2 \leq c_1$ .

**Case 2:**  $c_1 \in A \setminus \{a\}$  and  $c_2 = b$ . If  $c_1 \leq c_2$ , Then  $c_1 \leq b$ . Since  $b \leq a, c_1 \leq b \leq a$ . Thus  $c_1 \leq a$ , a contradiction. If  $c_2 \leq c_1$ , then  $b \leq c_1$ . Since  $a \leq b, a \leq b \leq c_1$ . So  $a \leq c_1$ , a contradiction.

**Case 3:**  $c_2 \in A \setminus \{a\}$  and  $c_1 = b$ . If  $c_1 \leq c_2$ , then  $b \leq c_2$ . Since  $a \leq b, a \leq b \leq c_2$ . Hence  $a \leq c_2$ , a contradiction. If  $c_2 \leq c_1$ , then  $c_2 \leq b$ . Since  $b \leq a, c_2 \leq b \leq a$ . Thus  $c_2 \leq a$ , a contradiction.

**Case 4:**  $c_1 = b$  and  $c_2 = b$ . This is impossible.

Thus  $B$  satisfies (1) and (2) in Theorem 3.1. Therefore  $B$  is a two-sided base of  $S$ .

**Corollary 3.3.** Let  $A$  be a two-sided base of an ordered  $G$ -semigroup  $(S, G, \leq)$ , and let  $a \in A$ . If  $I(x) = I(a)$  for some  $x \in S, x \neq a$ , then  $x$  belongs to two-sided base of  $S$ , which is different from  $A$ .

**Theorem 3.4.** Let  $A$  and  $B$  be any two-sided bases of an ordered  $G$ -semigroup  $(S, G, \leq)$ . Then  $A$  and  $B$  have the same cardinality.

**Proof.** Let  $a \in A$ . Since  $B$  is a two-sided base of  $S$ , by Theorem 3.1.(1), there exists an element  $b \in B$  such that  $a \leq b$ . Since  $A$  is a two-sided base of  $S$ , by Theorem 3.1.(1), there exists  $a^* \in A$  such that  $b \leq a^*$ . So  $a \leq b \leq a^*$ , i.e.,  $a \leq a^*$ . By Theorem 3.1.(2),  $a = a^*$ . Hence  $I(a) = I(b)$ . Define a mapping

$$j : A \rightarrow B \text{ by } j(a) = b \text{ for all } a \in A.$$

To show that  $j$  is well-defined, let  $a_1, a_2 \in A$  be such that  $a_1 = a_2, j(a_1) = b_1$  and  $j(a_2) = b_2$  for some



$b_1, b_2 \hat{=} B$ . Then  $I(a_1) = I(b_1)$  and  $I(a_2) = I(b_2)$ . Since  $a_1 = a_2$ ,  $I(a_1) = I(a_2)$ . Hence  $I(a_1) = I(a_2) = I(b_1) = I(b_2)$ , i.e.,  $b_1 \underline{\rho}_I b_2$  and  $b_2 \underline{\rho}_I b_1$ . By Theorem 3.1.(2),  $b_1 = b_2$ . Thus  $j(a_1) = j(a_2)$ . Therefore,  $j$  is well-defined. We will show that  $j$  is one- one. Let  $a_1, a_2 \hat{=} A$  be such that  $j(a_1) = j(a_2)$ . Since  $j(a_1) = j(a_2)$ ,  $j(a_1) = j(a_2) = b$  for some  $b \hat{=} B$ . So  $I(a_2) = I(a_1) = I(b)$ . Since  $I(a_2) = I(a_1)$ ,  $a_1 \underline{\rho}_I a_2$  and  $a_2 \underline{\rho}_I a_1$ . This implies  $a_1 = a_2$ . Therefore,  $j$  is one-one. We will show that  $j$  is onto. Let  $b \hat{=} B$ . Since  $A$  is a two-sided base of  $s$ , by Theorem 3.1.(1), there exists an element  $a \hat{=} A$  such that  $b \underline{\rho}_I a$ . Since  $B$  is a two-sided base of  $s$ , by Theorem 3.1.(1), there exists an element  $b^* \hat{=} B$  such that  $a \underline{\rho}_I b^*$ . So  $b \underline{\rho}_I a \underline{\rho}_I b^*$ , i.e.,  $b \underline{\rho}_I b^*$ . This implies  $b = b^*$ . Hence  $I(a) = I(b)$ . Thus  $j(a) = b$ . Therefore,  $j$  is onto. This completes the proof.

If a two- sided base  $A$  of an ordered  $G$ - semigroup  $(s, G, \xi)$  is a  $G$ - ideal of  $s$ , then  $s = (A \hat{=} sGA \hat{=} AGs \hat{=} sGAgs \hat{=} ] A ] = A$ . Hence  $s = A$ . The converse statement is obvious. Then we conclude that.

**Remark 3.5.** It is observed that a two-sided base  $A$  of an ordered  $G$ -semigroup  $(s, G, \xi)$  is a two-sided  $G$ -ideal of  $s$  if and only if  $A = s$ .

In Example 2.2., it is easy to see that  $\{d\}$  is a two-sided bases of  $s$ , but it is not a  $G$ -subsemigroup of  $s$ . This show that a two-sided bases of an ordered  $G$ -semigroup need not to be a  $G$ -subsemigroup in (Niovi, 2018). A non-empty subset  $A$  of  $s$  is called a idempotent if  $A = (AGA]$  or  $a = aga$  for all  $a \hat{=} A$  and  $g \hat{=} G$ . The following theorem gives necessary and sufficient conditions of a two-sided bases of  $s$  to be a  $G$ -subsemigroup  $s$ .

**Theorem 3.6.** A two-sided base  $A$  of an ordered  $G$ -semigroup  $(s, G, \xi)$  is a  $G$ -subsemigroup if and only if  $A = \{a\}$  with  $aga = a$  for all  $g \hat{=} G$ .

**Proof.** Assume that  $A$  of a  $G$ -subsemigroup  $s$ . Let  $a, b \hat{=} A$  and  $g \hat{=} G$ . Since  $A$  is a  $G$ -subsemigroup of  $s$ ,  $agb \hat{=} A$ . Setting  $agb = c$ ; thus  $c \hat{=} sGb \hat{=} sGb \hat{=} bGs \hat{=} sGbGB \hat{=} (sGb \hat{=} bGs \hat{=} sGbGB]$ . By Lemma 2.5.,  $c = b$ . So  $agb = b$ . Similarly,  $c \hat{=} aGS \hat{=} sGa \hat{=} aGS \hat{=} sGaGS \hat{=} (sGa \hat{=} aGS \hat{=} sGaGS]$ . By Lemma 2.5.,  $c = a$ . So  $agb = a$ . We have  $a = b$ . Therefore,  $A = \{a\}$  with  $aga = a$  for all  $a \hat{=} A$  and  $g \hat{=} G$ . The converse statement is clear.

**Notation.** The union of all two-sided bases of an ordered  $G$ -semigroup  $(s, G, \xi)$  is denoted by  $R$ .

**Theorem 3.7.** Let  $(s, G, \xi)$  be an ordered  $G$ -semigroup. Then  $s \setminus R$  is either empty set or a  $G$ -ideal of  $s$ .

**Proof.** Assume that  $s \setminus R \neq \emptyset$ . We will show that  $s \setminus R$  is a  $G$ -ideal of  $s$ . Let  $a \hat{=} s \setminus R$ ,  $x \hat{=} s$  and  $g \hat{=} G$ . To show that  $xga \hat{=} s \setminus R$  and  $agx \hat{=} s \setminus R$ . Suppose that  $xga \notin s \setminus R$ . Then  $xga \hat{=} R$ . Hence  $xya \hat{=} A$  for some a two-sided base  $A$  of  $s$ . We set  $b = xga$  for some  $b \hat{=} A$ . Then  $b \hat{=} sGa$ . By  $b \hat{=} sGa \hat{=} a \hat{=} sGa \hat{=} aGS \hat{=} sGaGS \hat{=} (a \hat{=} sGa \hat{=} aGS \hat{=} sGaGS]$  =  $I(a)$ , it follow that  $I(b) \hat{=} I(a)$ . Next we will show that  $I(b) \hat{=} I(a)$ .



Suppose that  $I(b) = I(a)$ . Since  $a \in S \setminus R$  and  $b \in A$ ,  $a \neq b$ . Since  $I(b) = I(a)$  and Corollary 3.3, we conclude that  $a \in R$ . This is a contradiction. Thus  $I(b) \not\subseteq I(a)$ , i.e.,  $b \notin I(a)$ . Since  $A$  is a two-sided base of  $S$  and  $a \in S \setminus R$ , by Theorem 3.1.(1), there exists  $d \in A$  such that  $a \leq_I d$ . Since  $b \notin I(a)$ ,  $a \leq_I d$ ,  $b \leq_I d$ . This is a contradiction to the condition (2) of Theorem 3.1., so we have  $xga \in S \setminus R$ . Similarly, to show that  $agx \in S \setminus R$ . Suppose that  $agx \in R$ , then  $agx \in A$  for some a two-sided base  $A$  of  $S$ . Let  $agx = c$  for some  $c \in A$ . Then  $c \in aGS$ . By  $c \in aGS \implies a \in Sga \in aGS \in SgaGS \implies (a \in Sga \in aGS \in SgaGS) = I(a)$ , it follow that  $I(c) \subseteq I(a)$ . Next, we will show that  $I(c) \not\subseteq I(a)$ . Suppose that  $I(c) = I(a)$ . Since  $a \in S \setminus R$  and  $c \in A$ ,  $a \neq c$ . Since  $I(c) = I(a)$  and Corollary 3.3., we conclude that  $a \in R$ . This is a contradiction. Thus  $I(c) \not\subseteq I(a)$ , i.e.,  $c \notin I(a)$ . Since  $A$  is a two-sided base of  $S$  and  $a \in S \setminus R$ , by Theorem 3.1.(1), there exists  $e \in A$  such that  $a \leq_I e$ . Since  $c \notin I(a)$ ,  $a \leq_I e$ ,  $c \leq_I e$ . This is a contradiction to the condition (2) of Theorem 3.1., so we have  $agx \in S \setminus R$ . Let  $x \in S \setminus R$ ,  $y \in S$  such that  $y \leq x$ . Next we will show that  $y \in S \setminus R$ . Suppose that  $y \in R$ , then  $y \in A$  for some a two-sided base  $A$  of  $S$ . Since  $A$  is a two-sided bases of  $S$ , by Theorem 3.1.(1) there exists an element  $z \in A$  such that  $x \leq_I z$ . Since  $y \leq x$ ,  $y \leq_I x$ . So we have  $y \leq_I z$ . This is a contradiction. Therefore  $y \notin R$  then  $y \in S \setminus R$ . Hence  $S \setminus R$  is a  $G$ -ideal of  $S$ .

**Notation.** Let  $M^*$  be a proper  $G$ -ideal of an ordered  $G$ -semigroup  $(S, G, \leq)$  containing every proper  $G$ -ideal of  $S$ .

**Theorem 3.8.** Let  $(S, G, \leq)$  be an ordered  $G$ -semigroup and  $\mathcal{A} \subseteq R \subseteq S$ . The following statements are equivalent:

- (1)  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ ;
- (2) For every element  $a \in R$ ,  $R \subseteq I(a)$ ;
- (3)  $S \setminus R = M^*$ ;
- (4) Every two-sided base of  $S$  is a one-element base.

**Proof.** (1)  $\implies$  (2). Assume that  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ . Let  $a \in R$ . Suppose that  $R \not\subseteq I(a)$ . Since  $R \not\subseteq I(a)$ , there exists  $x \in R$  such that  $x \notin I(a)$ . So we have  $x \in S \setminus R$ . Since  $x \notin I(a)$ ,  $x \in S \setminus R$  and  $x \in S$ , we have  $(S \setminus R) \cap I(a) \neq S$ . Thus  $(S \setminus R) \cap I(a)$  is a proper  $G$ -ideal of  $S$ . Hence  $S \setminus R \subseteq (S \setminus R) \cap I(a)$ . This contradicts to the maximality of  $S \setminus R$ .

Conversely, assume that for every element  $a \in R$ ,  $R \subseteq I(a)$ . We will show that  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ . Since  $a \in R$ ,  $a \in S \setminus R$ . Hence  $S \setminus R \subseteq S$ . Since  $R \subseteq S$ ,  $S \setminus R \subseteq \mathcal{A}$ . By Theorem 3.7.,  $S \setminus R$  is a proper  $G$ -ideal of  $S$ . Suppose that  $M$  is a proper  $G$ -ideal of  $S$  such that  $S \setminus R \subseteq M \subseteq S$ . Since  $S \setminus R \subseteq M$ , there exists  $x \in M$  such that  $x \in S \setminus R$ , i.e.,  $x \in R$ . Then  $x \in M \cap R$ . So  $M \cap R \subseteq \mathcal{A}$ . Let  $c \in M \cap R$ . Then  $c \in M$  and  $c \in R$ . Since  $c \in M$ ,  $Sgc \in SGM \subseteq M$ ,  $cGS \subseteq MGS \subseteq M$  and  $SgcGS \subseteq SGMGS \subseteq M$ . Then





$I(c) = (c \in S G c \in c G S \in S G c G S) \cap M$ . Since  $c \in R$ , by assumption we have  $R \subseteq I(c)$ . Hence  $S = (S \setminus R) \cup R \subseteq (S \setminus R) \cup I(c) \subseteq M \subseteq S$ . Thus  $M = S$ . This is a contradiction. Therefore  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ .

(3)  $\Rightarrow$  (4). Assume that  $S \setminus R = M^*$ . Since  $S \setminus R = M^*$ ,  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ . By (1)  $\Rightarrow$  (2), for every  $a \in R, R \subseteq I(a)$ . First, we will show that for every  $a \in R, S \setminus R \subseteq I(a)$ . Suppose that  $S \setminus R \not\subseteq I(a)$  for some  $a \in R$ . Then  $I(a) \subsetneq S$ . Hence  $I(a)$  is a proper  $G$ -ideal of  $S$ . Thus  $I(a) \subseteq M^* = S \setminus R$ . Then  $I(a) \subseteq S \setminus R$ . Since  $a \in I(a), a \in S \setminus R$ , i.e.,  $a \notin R$ . This is a contradiction. Thus  $S \setminus R \subseteq I(a)$  for every  $a \in R$ . Since  $S \setminus R \subseteq I(a)$  and  $R \subseteq I(a)$  for every  $a \in R$ , it follows that  $S = (S \setminus R) \cup R \subseteq I(a) \cup I(a) = I(a) \subseteq S$ . So  $S = I(a)$  for every  $a \in R$ . Therefore,  $\{a\}$  is a two-sided base of  $S$ . Let  $A$  be a two-sided of  $S$ . We will show that  $a = b$  for all  $a, b \in A$ . Suppose that there exist  $a, b \in A$  such that  $a \neq b$ . Since  $A$  is a two-sided base of  $S, A \subseteq R$ . This is,  $a \in R$ . So  $S = I(a)$ . Since  $b \in S = I(a)$  and  $b \neq a, b \in (S G a \in a G S \in S G a G S)$ . By Lemma 2.5.,  $a = b$ . This is a contradiction. Therefore, every two-sided base of  $S$  is a one element base.

Conversely, assume that every two-sided base of  $S$  is a one element base. Then  $S = I(a)$  for all  $a \in R$ . We will show that  $S \setminus R = M^*$ . The statement that  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$  follows from the proof (1)  $\Rightarrow$  (2). Let  $M$  be a  $G$ -ideal of  $S$  such that  $M$  is not contained in  $S \setminus R$ . Then  $R \cap M \neq \emptyset$ . Let  $a \in R \cap M$ . Hence  $a \in R$  and  $a \in M$ . So  $S G a \in S G M \subseteq M, a G S \in M G S \subseteq M$  and  $S G a G S \subseteq S G M G S$ . So we have  $I(a) = (a \in S G a \in a G S \in S G a G S) \cap M$ . Hence  $S = I(a) \subseteq M \subseteq S$ . Thus  $M = S$ . Therefore  $S \setminus R = M^*$ .

(1)  $\Rightarrow$  (3). Assume that  $S \setminus R$  is a maximal proper  $G$ -ideal of  $S$ . We will show that  $S \setminus R = M^*$ . Since  $S \setminus R$  is a proper  $G$ -ideal of  $S, S \setminus R \subseteq M^* \subseteq S$ . By assumption,  $S \setminus R = M^*$  or  $S = M^*$ . Since  $S \neq M^*$ , so we have  $S \setminus R = M^*$ . The converse statement is obvious.

## Discussion

In this research, we investigated the notion of ordered  $G$ -semigroup containing two-sided bases. We proved that a non-empty subset  $A$  of an ordered  $G$ -semigroup  $(S, G, \leq)$  is a two-sided base of  $S$  if and only if  $A$  satisfies the following two conditions (1) for any  $x \in S$  there exists  $a \in A$  such that  $x \leq_a a$ ; (2) for any  $a, b \in A$ , if  $a \leq b$ , then neither  $a \leq b$  nor  $b \leq a$ . Also, we showed that if  $A$  and  $B$  be any two-sided bases of an ordered  $G$ -semigroup  $(S, G, \leq)$ . Then  $A$  and  $B$  have the same cardinality.

## Conclusions

In this research, we have result in ordered  $G$ -semigroup that are analogously in  $G$ -semigroup considered by T. Changpas and P. Kummoon in 2018.



## References

- Abul Basar, M.Y. Abbasi and Sabahat Ali Khan. (2017). Some properties of covered  $G$ -ideal in po- $G$ -semigroups. *International Journal of Pure and Applied Mathematics*, 115(2), 345-352.
- Chinnadurai, V., & Arulmozhi, K. (2018). Characterization of bipolar fuzzy ideal in ordered gamma semigroup. *Journal of the International Mathematical Virtual Institute*, 8, 141-156.
- Fabrics, I. (1975). Two sided bases of semigroups. *Matematicky casopis*, 25(2), 173-178.
- Iampan, A. (2009). Characterizing Ordered Bi-Ideal in Ordered  $G$ -Semigroups. *International Journal of Mathematical Sciences and Informatics*, 4(1), 17-25.
- Kostaq Hila & Edmond Pisha. (2006). Characterizations on ordered  $G$ -semigroups. *International Journal of Pure and Applied Mathematics*, 28(3), 423-439.
- Niovi Kehayopulu. (2011). On regular duo po- $G$ -semigroup. *Mathematical Institute Slovak Academy of Sciences*, 61(6), 871-884.
- Kwon Y.I. and Lee S.K. (1998). On weakly prime ideals of ordered  $G$ -semigroup. *Comm. Korean Math. Soc.*, 13(2), 251-256.
- Niovi Kehayopulu. (2017). On intra-regular ordered  $G$ -semigroup. *European Journal of Pure and Applied Mathematics*, 10(4), 620-630.
- Sen, M. K. (1981). On  $G$ -semigroup. *Lecture Notes in Pure and Appl. Math.*, 91, 301-308.
- Subrahmanyaswara Rao Seetamraju, V.B., A. Anjaneyulu, A., & Madhusudana Rao, D. (2012). Po- $G$ -ideals in Po- $G$ -semigroup. *IOSR Journal of Mathematics*, 1(6), 39-51.
- Thawat Chang phas & Pisit Kummoon. (2018). On  $G$ -Semigroup Containing Two-sided bases. *KKU Sci. J.*, 46(1), 154-161.



Thawhat Changphas & Pisit Kummoon. (2018). On left and right bases of a  $\mathcal{G}$ -semigroup. *International Journal of Pure and Applied Mathematics*, 118(1), 125-135.